

DOI:10.11931/guihaia.gxzw201807002

苹果菊属，东南亚大陆菊科一新归化属

沈佳豪^{1,2}, 牛帼豪^{1,2}, 张国进^{1,2}, 付志玺³, INTA Angkhana⁴, 高天刚^{1,2*}

(1. 中国科学院植物研究所, 系统与进化植物学国家重点实验室, 中国 北京 100093; 2. 中国科学院大学, 中国 北京 100049; 3. 四川师范大学, 生命科学学院, 中国 成都 610101; 4. 清迈大学, 理学院, 生物学部, 泰国 清迈 50200)

摘要: 报道了在泰国北部和中国东南部的菊科一个新归化属——苹果菊属 (*Centratherum* Cass.)。依据所采集的标本, 详细描述了该属在东南亚大陆的唯一代表亚种苹果菊 [*Centratherum punctatum* Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li] 并配以插图。苹果菊与其所在属的其它物种之区别在于该亚种叶片菱形至椭圆形, 叶缘具不规则锯齿, 先端微尖, 总苞片基部硬化, 先端无芒。文中提供了苹果菊的分布图, 并讨论了其可能的传播途径。

关键词: 苹果菊属, 菊科, 斑鸠菊族, 归化属, 东南亚大陆, 命名法

中图分类号: Q949

文献标识码: A

Centratherum, a newly naturalized genus of the family

Asteraceae from mainland of Southeast Asia

SHEN Jiahao^{1,2}, NIU Guohao^{1,2}, ZHANG Guojin^{1,2}, FU Zhixi³,
INTA Angkhana⁴, GAO Tiangang^{1,2*}

(1. State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China; 2. University of the Chinese Academy of Sciences, Beijing 100049, China; 3. College of Life Sciences, Sichuan Normal University, Chengdu 610101, China; 4. Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand)

Abstract: *Centratherum* Cass., a newly naturalized genus of the family Asteraceae from Southeast Asia, was recorded in northern Thailand and southwestern China. *Centratherum punctatum* Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li, the only one representative subspecies of this genus from mainland of Southeast Asia, is described and illustrated based on the specimens. *C. punctatum* subsp. *fruticosum* can be distinguished from the other taxa of this genus by its leaf blade rhombic to elliptic, leaf margins irregularly serrate, leaf apex broadly

基金项目: 国家自然科学基金(31570204, 31270237, J1310002); 国家科技部科技基础性工作专项(2014FY210300, 2013FY112100) [Supported by the National Natural Science Foundation of China (31570204, 31270237, J1310002); Science and Technology Basic Work of the Ministry of Science and Technology of China (2014FY210300, 2013FY112100)].

作者简介: 沈佳豪(1994-), 男, 浙江湖州人, 硕士研究生, 主要从事植物分类学研究, (E-mail) shenjiahao17@mails.ucas.ac.cn。

***通信作者:** 高天刚, 博士, 副研究员, 主要从事植物分类和系统进化方面的研究, (E-mail) gaotg@ibcas.ac.cn。

acute, and phyllaries indurate at base and not awned. Its distribution map is provided and the possible dispersal pathway is discussed.

Keywords: *Centratherum*, Asteraceae, Vernoniaeae, naturalized genus, mainland of Southeast Asia, nomenclature

In a field trip to Thailand in May 2014, we collected an unknown taxon of the tribe Vernoniaeae of Asteraceae. Similar specimens were also collected in an expedition to Yunnan, China in April 2017. We identified these specimens as *Centratherum punctatum* Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li, a genus and a species not previously recorded in the mainland of Southeast Asia. *Centratherum* Cass. is a tropical genus distributed in tropical America, Australia and Philippines (Kirkman, 1981). *C. punctatum* subsp. *fruticosum* is native to Philippines and can be distinguished from the other taxa of this genus by its leaf blade rhombic to elliptic, leaf margins irregularly serrate, leaf apex broadly acute, and phyllaries indurate at base and not awned (Kirkman, 1981; Chen et al, 1999).

Based on the specimens collected during the flowering phase, we provided the description as below.

Centratherum punctatum Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li in *Taiwania* 44(2): 300. 1999 (Figs. 2–4).

≡ *Centratherum fruticosum* S. Vidal, *Rev. Pl. Vasc. Filip.* 159. 1886; Elmer in *Leafl. Philip. Bot.* 1: 88. 1906.

Type:—PHILIPPINES. Distr. Lepanto (Mt. Province), 1876, *S. Vidal 1502* (syntypes: A photo! [No. A00004641], L photo! [No. L.3034111], MA photo! [No. MA728008]).

= *Centratherum punctatum* Cass. subsp. *fruticosum* (Elmer) K. Kirkman in *Rhodora* 83: 20. 1981, nom. inval.

Type:—PHILIPPINES. Prov. Batangas, Luzon, 1841, *H. Cuming 1556* (lectotype (designated by Kirkman 1981): G photo! [No. G00223244]; isolectotypes: E photo! [No. E00417342], K photo! [No. K000796695], LE photo! [No. LE00017869, LE00017870]).

Perennial herbs or subshrubs, up to 1–1.4 m tall, branched (Fig. 1: A). Stem woody glabrous at base, young shoot hirsute, ridged. Leaves crowded, short internodes often appear fascicled, leaf blade elliptic-obovate to rhombic, 1–8.3 × 0.5–3.9 cm, broadly acute at apex, attenuate to truncate at base, margin irregularly crenate-serrate, uniseriate and T-shaped trichomes on both surfaces; petiole up to 2 cm, winged, indistinct in small blades. Capitula terminal or terminal on axillary branches, 2–2.8 cm in diam., 1.8–2 cm tall; peduncle 3–11 cm long, pubescent, receptacle naked; involucre campanulate to sub-globose, ca. 1 cm tall, base with leafy whorled bracts; phyllaries imbricate in 5–7 series, minutely pubescent, 3-veined, inner phyllaries linear-oblongate to fiddle-shaped, 8 × 2.5–2.7 mm, apex mucronate to cuspidate, margin scarious, outer phyllaries triangular, 5–7 × 1–1.2 mm, apex spine or acuminate, base indurate (Fig. 1: B). Capitulum with many florets, hermaphrodite, regular. Corolla tubular, reddish-purple, 8–16 mm long, outer part with glandular hairs, lobes 5, linear-lanceolate (Fig. 1: C). Style white, style-branches filiform, flushed with purple and hispidulous (Fig. 1: D). Stamens 5, white; anthers linear, 3 × 0.5 mm, appendant lanceolate, base calcarate (Fig. 1: E). Achene obconic-cylindrical, ca. 2.5 × 1.5 mm, black, glabrous, with 10 ribs. Pappus of 6–10 bristles, deciduous, straw-colored, 2–3 × 0.2 mm.

Phenology: Flowering and fruiting throughout the years.

Specimens examined: THAILAND. Chiang Rai: Maesai to Nan, alt. 364 m, 99°51'32" E,

20°12'17" N, 14 May 2014, *Tian-Gang Gao & Guo-Jin Zhang 5807* (PE). CHINA. Yunnan: Dehong Autonomous Prefecture, Ruili City, Wanding Town, Huihuan Village, alt. 935 m, 98°2'40" E, 24°7'4" N, 1 April 2017, *Jin-Pu Wei et al. RL0832* (HCNGB). (Fig. 2)

In Asia, *Centratherum punctatum* Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li was recorded in Taiwan (Chen et al, 1999) and West Bengal (Chowdhury & Das, 2012). But it has never been recorded in the mainland of Southeast Asia. Besides the present distribution sites (southwestern China and northern Thailand), it could also possibly be discovered in the nearby places (like Myanmar) in the future. It may be dispersed to the continent of Southeast Asia by flower trade, as marketing information about it can be obtained in two big business websites of China (Jingdong, 2018; Taobao, 2018). Hitherto, *C. punctatum* subsp. *fruticosum* has not been reported as an invasive species. However, based on our field observations, it could produce an immense number of mature achenes (ca. 150) in one capitulum. In addition, because of the vegetational and climatic similarities between mainland of southeast Asia and origin distribution (Zhang et al, 2017), close attention should be paid to its potential dispersal.

Note: *Centratherum fruticosum* was validly published by Vidal (1886) with indicating a single gathering (*Vidal 1502*). Quisumbing (1958, in sched.) labeled one specimen of this gathering (MA [No. MA728008]) as “type”, the other (A [No. A00004641]) as “isotype”, but no effective publication was found. According to Art 7.9 and 9.5 of ICN (McNeill et al, 2012), these three specimens from a single gathering should be treated as syntypes. In 1906, Elmer used the name *C. fruticosum* S. Vidal by giving a full and direct reference, and cited the specimens *Cuming 1556* and *Elmer 5956*. In 1981 Kirkman established the new combination *C. punctatum* Cass. subsp. *fruticosum* (Elmer) K. Kirkman based on the incorrect author and reference citation. In addition, Kirkman designated *Cuming 1556* (G [No. G00223244]) as lectotype of *C. punctatum* Cass. subsp. *fruticosum* (Elmer) K. Kirkman and described it only in English. According to the Art 39.1, 41.5 and 41.8(d) of ICN (McNeill et al, 2012), as a new taxon name, *C. punctatum* subsp. *fruticosum* K. Kirkman was invalidly published because of no Latin description or diagnosis. Hence, the new combination *C. punctatum* subsp. *fruticosum* (Elmer) K. Kirkman was invalidly published because of the indirect reference. In 1999 Chen et al. cited Kirkman’s publication but clearly indicated the correct basionym *C. fruticosum* S. Vidal with a full and direct reference. Hence, *C. punctatum* Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li was the valid name. The result is coincident with that of Chowdhury & Das (2012).


Acknowledgements

We are indebted to the curators of the herbarium PE and HCNGB for access to specimens.

参考文献:

- CHEN SH, WU MJ, LI SM, 1999. *Centratherum punctatum* Cass. ssp. *fruticosum*, a newly naturalized sunflower species in Taiwan[J]. Taiwania, 44: 299–305.
- CHOWDHURY M, DAS AP, 2012. Record of naturalization of *Centratherum punctatum* Cassini ssp. *fruticosum* (S. Vidal) K. Kirkman (Asteraceae) in India with a note on its nomenclature[J]. J Bot Soc Bengal, 66: 69–71.
- ELMER ADE, 1906. Manual of the Philippine Compositae[J]. Leaflet Philip Bot, 1: 83–186.
- JINGDONG, 2018 [2018-07-01]. [EB/OL]. <https://item.jd.com/24598283886.html>.

- KIRKMAN LK, 1981. Taxonomic revision of *Centratherum* and *Phyllocephalum* (Compositae: Vernonieae)[J]. Rhodora, 83: 1–24.
- MCNEILL J, BARRIE FR, BUCK WR, et al, 2012. International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011 (Regnum Vegetabile 154)[M]. Königstein: Koeltz Scientific Books.
- TAOBAO, 2018 [2018-07-01]. [EB/OL]. <http://m.tb.cn/h.WuTJpcx>.
- VIDAL S, 1886. Revision de plantas vasculares Filipinas[M]. Manila: Establecimiento Tipo-Litográfico de M. Perze: 159.
- ZHANG X, WU S, YAN X, et al, 2017. A global classification of vegetation based on NDVI, rainfall and temperature[J]. Int J Climatol, 37: 2318–2324.



注：A. 习性；B. 头状花序；C. 花冠；D. 花柱分枝；E. 花药。

Note: A. Habit; B. Capitulum; C. Corolla; D. Style branches; E. Anther.

图 1 苹果蓟

Fig. 1 *Centratherum punctatum* Cass. subsp. *fruticosum* (S. Vidal) K. Kirkman ex Shih H. Chen, M. J. Wu & S. M. Li




图 2 苹果薊在东南亚大陆的分布

Fig. 2 Distribution of *Centratherum punctatum* subsp. *fruticosum* in mainland of Southeast Asia